

(12) **UK Patent Application** (19) **GB** (11) **2 176 786 A**
(43) Application published 7 Jan 1987

(21) Application No 8614798

(22) Date of filing 18 Jun 1986

(30) Priority data

(31) 60/135718

(32) 20 Jun 1985

(33) JP

(71) Applicant

Sumitomo Chemical Company Limited

(Incorporated in Japan)

15 Kitahama 5-chome, Higashi-ku, Osaka shi, Osaka
541, Japan

(72) Inventors

Ayumu Inoue

Takeaki Umemura

(74) Agent and/or Address for Service

J. A. Kemp & Co.

14 South Square, Gray's Inn, London WC1R 5EU

(51) INT CL⁴

C07C 57/58 87/28

(52) Domestic classification (Edition I):

C2C 220 226 227 22Y 239 29X 29Y 30Y 313 31Y 321

32Y 338 366 367 37X 490 618 620 628 650 658 699 776

778 779 BU NN VN

(56) Documents cited

GB A 2014137

GB A 2013670

EP A1 0107972

(58) Field of search

C2C

(54) **A method for optical resolution of phenylacetic acid derivative**

(57) Alpha-isopropyl-p-chlorophenylacetic acid, which is an acid moiety of the pyrethroidal ester pesticides is subjected to optical resolution using optically active alpha-phenyl-beta-p-tolyethylamine or optically active alpha-phenylethylamine, as the optical resolution agent. The reaction of the acid with the amine is carried out in a hydrophobic organic solvent, such as aliphatic and aromatic hydrocarbons, in the presence of water, until the desired salt of the acid with the amine is separated therein through selective crystallization.

GB 2 176 786 A

SPECIFICATION

A method for optical resolution of phenylacetic acid derivative

5 This invention relates to a method for the resolution of alpha-isopropyl-p-chlorophenylacetic acid (ICPA). More particularly, it relates to a method for the optical resolution of ICPA using optically active alpha-phenyl-beta-p-tolyethylamine (PTE) or optically active alpha-phenylethylamine (PEA) in a hydrophobic organic solvent in the presence of water. 5

10 It has already been known that a group of alpha-substituted phenylacetic acid esters having a chemical structure greatly different from that of the conventional pyrethroidal insecticides has a strong insecticidal activity against various harmful insects (United Kingdom Patent No. 1439615), and, among them, the esters of ICPA are particularly excellent in its efficacy and economy. 10

Furthermore, as to the insecticidal efficacy of the esters of the (+)- and (-)-carboxylic acids obtained by the optical resolution of its alpha-substituted phenylacetic acid moiety, it has been 15 known that the ester of the (+)-acid exhibits about twice a strong insecticidal effect as compared with that of the ester of the (±)-acid, while the ester of (-)-acid is almost ineffective. 15

The present inventors have extensively studied to develop the optical resolution method which is expectedly more efficient from the commercial viewpoint, and have discovered that, in the method for optically resolving ICPA using optically active PTE or optically active PEA, (+)-ICPA 20 of a high optical purity could be obtained with satisfactory yield by using a hydrophobic solvent and operating in the presence of water. The present invention has been accomplished according to such findings. 20

Thus this invention provides a method for the optical resolution of ICPA, using optically active PTE or optically active PEA as the optical resolution agent, which method is characterized by 25 that the reaction of the acid with the amine is carried out in a hydrophobic organic solvent in the presence of water, until one of the salts of the optically active ICPA with the amine is separated therein through selective crystallization. 25

According to the method of the present invention, one of the salt of the optically active ICPA with the amine can be obtained in high purity and in the form of readily filterable and (+)-ICPA 30 of a high optical purity can be obtained without any additional purifying process such as recrystallization procedure. 30

As a secondary effect of the method of this invention, it is advantageous that the manufacturing and purifying processes of ICPA, if conducted in the method mentioned, for example, in the Japanese Unexamined Patent Publications Nos. 5350/1975 and 25535/1978, can be directly 35 connected to the resolution process of this invention, wherein the ICPA solution in a hydrophobic organic solvent can be used as it is. Namely, as for the method for synthesizing ICPA, for example, as mentioned in the Japanese Unexamined Patent Publication No. 5350/1975, hydrolysis of alpha-isopropyl-p-chlorophenylacetonitrile has been known. And in the Publication No. 25535/1978, a method for purifying ICPA using benzene, toluene, xylene, chlorobenzene, hexane, heptane, or any other aliphatic or aromatic organic solvent has been disclosed. In conducting 40 the method of this invention, it is not necessary to isolate the ICPA, and the solution of ICPA in an aliphatic or aromatic organic solvent used in the above refining method directly be employed for the present optical resolution process. 40

Furthermore, since a hydrophobic organic solvent is used according to this method, it is 45 extremely easy to recover the solvent. In this regard, this method is quite advantageous for conducting the optical resolution of ICPA in an industrial scale, in particular. 45

Thus, this invention relates to a method for the optical resolution of ICPA, which produces (+)-ICPA of a high optical purity at a high efficiency with constant yield, even when carried out in an industrial scale, together with various other commercial advantages. 50

The method of this invention will be further described below. 50

Examples of the hydrophobic organic solvents used in this invention include benzene, toluene, xylene and other aromatic hydrocarbons, and hexane, heptane, octane and other aliphatic hydrocarbons.

In this method, water is used in an amount ranging from 3 to 10 wt.%, based upon the 55 amount of ICPA. 55

As the optically active PTE or PEA, (+)-PTE or (-)-PEA is preferable, and its amount is preferably in a range of 0.35 to 0.65 mol per mol of (±) ICPA. The optical purity of these optical resolution agents may preferably be 92.0% e.e. or more.

Among these optical resolution agents, it is preferable to use (+)-PTE in an amount ranging 60 from 0.45 to 0.65 mol per mol of (±) ICPA, from the viewpoint of optical purity of (+)-ICPA to be obtained. 60

The reaction period of time is not particularly limited, if it is more than 30 minutes. The amount of the organic solvent is preferably in a range from 1 to 10 times by weight, more preferably, 2 to 4 times by weight, that of the ICPA used.

65 The method of this invention is conducted, for example, in the following procedure. In the first 65

place, (\pm)-ICPA is caused to react with (+)-PTE or (-)-PEA in the solvent as mentioned above in the presence of appropriate amount of water. The temperature in this reaction is not limited, but it is desirable to keep the temperature at 40 to 150°C during or after the reaction, in order to obtain (+)-ICPA of a higher optical purity. While keeping the temperature at 40 to 150°C, it is not necessary that the salt produced is completely dissolved. After keeping at the temperature, the precipitated salt of (+)-ICPA with the amine, preferably by cooling, is separated from the mother liquor. At this time, the ICPA remaining in the mother liquor is in the (-)-form. The separating temperature is preferably 0 to 60°C, or more preferably 10 to 30°C.

The salt of optically active ICPA may directly be used for the manufacturing process of an ester of ICPA, but, more preferably, the salt may be once transformed into an optically active ICPA or its alkaline salt by an ordinary method using an aqueous acid, such as aqueous hydrochloric acid and aqueous sulfuric acid, or an aqueous alkali, such as aqueous sodium hydroxide and aqueous potassium hydroxide, and then derived to an ester of ICPA.

This invention will be described below in further details by referring to the examples, to which, however, this invention is not limited. Meanwhile the optical purity of ICPA shown in the following examples were determined by the method mentioned in Agric. Biol. Chem., 43, 2311 (1979), by Horiba et al., after decomposing the obtained salt of ICPA according to the ordinary method.

Examples 1-3

To 61.04 g of (\pm)-ICPA (chemical purity: 98.3%) was added 107.10 g of toluene, and the mixture was stirred to form a solution. Water was added thereto in an amount as described in Table 1. A solution of a described amount of (+)-PTE (optical purity: 95.0% e.e.) in 72.90 g of toluene was added to the above solution, and the mixed solution was heated up to 75°C. After keeping the temperature at 75°C for one hour, the solution was allowed to cool at a rate of about 1°C/5 minutes. After reaching 20°C, the solution was held for one hour at the same temperature. The precipitated crystals were isolated by filtration, washed with a small volume of toluene, and dried. The salt of (+)-ICPA with (+)-PTE was obtained in an amount shown in Table 1.

At the same time, the similar operation was conducted without adding water, as a reference example.

Table 1

Example No.	Amount of water used (g)	Amount of (+)-PTE used (g)	Obtained amount of (+)-ICPA-(+)-PTE salt		Optical purity of ICPA obtained (% e.e.)
			Yield (g)	Yield (%) (vs. amount of (\pm)-ICPA)	
Example 1	2.4	31.24	37.68	31.5	94.6
Example 2	6.0	31.24	37.29	31.2	94.4
Example 3	2.4	37.37	47.53	39.7	93.6
Reference example 1	0	31.24	39.78	33.3	89.8

To 35.00 g of the salt obtained in Example 1 in the above were added 60.00 g of toluene and 72.60 g of 5% aqueous sodium hydroxide solution, and the mixture was stirred at 40°C for 1 hour. Then the reaction mixture was separated to an aqueous layer and a toluene layer. The aqueous layer was extracted with toluene and the toluene layer was combined with the toluene layer obtained in the above. The combined toluene layer was concentrated to give 17.39 g of (+)-PTE. The aqueous layer was acidified with 15% aqueous sulfuric acid and extracted with toluene. The toluene layer was washed with water and concentrated to give 17.53 g of (+)-ICPA.

Example 4

To 61.04 g of (\pm)-ICPA (chemical purity: 98.3%) was added 107.10 g of toluene, and the mixture was stirred to make a solution. Then, 2.4 g of water was added thereto. A solution of 31.67 g of (+)-PTE (optical purity: 92.0% e.e.) in 72.90 g of toluene was added to the above solution, and the mixture was heated to 75°C. After keeping the temperature at 75°C for one hour, the mixed solution was allowed to cool at a rate of about 1°C/5 minutes down to 20°C. Then the solution was held for additional one hour at 20°C. The precipitated crystals were isolated by filtration, washed with a small volume of toluene, and dried, to obtain the salt of (+)-ICPA with (+)-PTE in an amount shown in Table 2.

At the same time, the similar operation was conducted without adding water, as a reference example.

Table 2

Example No.	Amount of water used (g)	Obtained amount of (+)-ICPA-(+)-PTE salt		Optical purity of ICPA obtained (% e.e.)
		Yield (g)	Yield (%) (vs. amount of (\pm)-ICPA)	
Example 4	2.4	37.06	31.0	92.0
Reference example 2	0	38.87	32.5	82.6

Example 5

To 61.04 g of (\pm)-ICPA (chemical purity: 98.3%) was added 107.10 g of toluene, and the mixture was stirred to make a solution. Then, 2.4 g of water was added thereto. A solution of 13.90 g of (-)-PEA (optical purity: 96.0% e.e.) in 72.90 g of toluene was added to the above solution, and the mixture was heated to 75°C. After keeping the temperature after 75°C for one hour, the mixed solution was allowed to cool at a rate of about 1°C/5 minutes down to 20°C. Then the solution was held for additional one hour at 20°C. The precipitated crystals were isolated by filtration, washed with a small volume of toluene, and dried, to obtain the salt of (+)-ICPA with (-)-PEA in an amount shown in Table 3.

At the same time, the similar operation was conducted without adding water, as a reference example.

Table 3

Example No.	Amount of water used (g)	Obtained amount of (+)-ICPA-(-)-PEA salt		Optical purity of ICPA obtained (% e.e.).
		Yield (g)	Yield (%) (vs. amount of (\pm) -ICPA)	
Example 5	2.4	33.44	35.5	87.0
Reference example 3	0	32.68	34.7	81.8

CLAIMS

1. A method for the optical resolution of racemic alpha-isopropyl-p-chlorophenylacetic acid, which comprises reacting the racemic acid with, as optical resolution agent, an amine which is optically active alpha-phenyl-beta-p-tolyethylamine or optically active alpha-phenylethylamine in a hydrophobic organic solvent in the presence of water, and separating a salt of optically active alpha-isopropyl-p-chlorophenylacetic acid with the amine through selective crystallization.
2. A method according to claim 1, wherein said amine is used in an amount of 0.35 to 0.65 mol per mol of alpha-isopropyl-p-chlorophenylacetic acid.
3. A method according to claim 1 or 2, wherein (+)-alpha-phenyl-beta-p-tolyethyl-amine or (-)-alpha phenylethylamine is used as the optical resolution agent.
4. A method for the optical resolution according to claim 3, wherein (+)-alpha-phenyl-beta-p-tolyethylamine is used in an amount of 0.45 to 0.65 mol per mol of alpha-isopropyl-p-chlorophenylacetic acid.
5. A method according to any one of the preceding claims wherein water is used in an amount of 3 to 10 wt.% based upon the amount of alpha-isopropyl-p-chlorophenylacetic acid.
6. A method according to any one of the preceding claims wherein the hydrophobic organic solvent is at least one aromatic hydrocarbon or aliphatic hydrocarbon.
7. A method according to any one of the preceding claims, wherein the solvent is used in an amount of 2 to 4 times by weight based upon the amount of alpha-isopropyl-p-chlorophenylacetic acid.
8. A method according to claim 1 substantially as described with reference to any one of the Examples.
9. Use of optically active alpha-isopropyl-p-chlorophenylacetic acid obtained, as such or in the form of an ester with the amine, by a method as claimed in any one of the preceding claims in the preparation of insecticidally active pyrethroids.

Printed in the United Kingdom for Her Majesty's Stationery Office, Dd 8818935, 1987, 4235.

Published at The Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from which copies may be obtained.